VOL. 550 NOS. 1+2 JOURNAL OF CHROMATOGRAPHY JULY 26, 1991

## CONTENTS

## HONOUR VOLUME ON THE OCCASION OF THE 60TH BIRTHDAY OF LLOYD R. SNYDER

Preface
by Cs. Horváth . . . . . 1

OPTIMIZATION

Computer-assisted retention prediction system for inorganic cyclic polyphosphates and its application to optimization of gradients in anion-exchange chromatography
by Y. Baba (Kobe, Japan) and G. Kura (Fukuoka, Japan) . . . . . 5

Multiple-criteria optimization
by S. N. Deming (Houston, TX, USA) . . . . . 15

Fully automatic high-performance liquid chromatographic optimization
by N. M. Djordjevic, F. Erni and B. Schreiber (Basle, Switzerland), E. P. Lankmayr and W. Wegscheider (Graz, Austria) and L. Jaufmann (Munich, Germany) . . . . . 27

Fast development of a robust high-performance liquid chromatographic method for *Ginkgo biloba* based on computer simulation
by I. Molnar (Berlin, Germany) and K. H. Gober and B. Christ (Köln, Germany) . . . . 39

Resolution enhancement of chromatographic data. Considerations in achieving super-resolution with the constrained iterative relaxation method
by M. R. Schure (Spring House, PA, USA) . . . . . 51

Computer-aided optimization of high-performance liquid chromatographic analysis of flavonoids from some species of the genus *Althaea*
by T. H. Dzido and E. Soczewiński (Lublin, Poland) and J. Gudej (Łódź, Poland) . . . . 71

Computer simulation of isocratic retentions of alkylketones using gradient data
by J. D. Stuart and D. D. Lisi (Storrs, CT, USA) . . . . . 77

Prediction of initial high-performance liquid chromatographic conditions for selectivity optimization in pharmaceutical analysis by an expert system approach
by G. Szepesi and K. Valkó (Budapest, Hungary) . . . . . 87

GAS CHROMATOGRAPHY

Measurement of solute dipolarity/polarizability and hydrogen bond acidity by inverse gas chromatography
by J. Li, Y. Zhang, A. J. Dallas and P. W. Carr (Minneapolis, MN, USA) . . . . . 101

Effect of stationary phase on predictions of the statistical model of overlap from gas chromatograms
by F. J. Oros and J. M. Davis (Carbondale, IL, USA) . . . . . 135

Gas chromatography of homologous esters. XXXIV. Alkyl borate and boronate esters
by J. K. Haken and F. Abraham (Kensington, Australia) . . . . . 155

Gas chromatographic study of interaction between vinyl chloride and poly(vinyl chloride)
by A. Koliadima, P. Agathonos and G. Karaiskakis (Patras, Greece) . . . . . 171

Homogeneous catalysis studied by reversed-flow gas chromatography
by B. V. Stolyarov (Leningrad, USSR) and N. A. Katsanos, P. Agathonos and J. Kapolos (Patras, Greece) . . . . . 181

Tandem mass spectrometry —an additional dimension of chromatography for the determination of biomarkers in fossil fuels (Review)
by R. P. Philp (Norman, OK, USA), J.-N. Oung (Taipei, Taiwan) and A. Galvez-Sinibaldi and J. Allen (Norman, OK, USA) . . . . . 193

Application of multivariate analysis to the selection of test solutes for characterizing stationary phase selectivity in gas chromatography
by T. O. Kollie and C. F. Poole (Detroit, MI, USA) . . . . . 213

Diurnal changes in plasma levels of 2-pyrrolidinone determined by isotope dilution mass spectrometry
by G. A. van der Berg, B. G. Wolthers, G. T. Nagel and F. A. J. Muskiet (Groningen, Netherlands), S. Banghman (Richmond, VA, USA) and R. A. de Zeeuw (Groningen, Netherlands) . . . . . 239

COLUMN LIQUID CHROMATOGRAPHY
*Theory*

Equilibrium elution chromatography. The column-origin boundary condition
by P. J. Karol (Pittsburgh, PA, USA) . . . . . 247

Expert system for repeatability testing of high-performance liquid chromatographic methods
by M. Mulholland and N. Walker (Cambridge, UK), F. Maris and H. Hindriks (Oss, Netherlands), L. Buydens (Nijmegen, Netherlands), T. Blaffert (Hamburg, Germany) and P. J. Schoenmakers (Eindhoven, Netherlands) . . . . . 257

Role of $\pi$–$\pi$ interaction in reversed-phase liquid chromatography
by G. Thévenon-Emeric, A. Tchapla and M. Martin (Palaiseau, France) . . . . . 267

Reversed-phase liquid chromatographic study of excess and absolute sorption isotherms of acetonitrile–water mixtures
by A. Alvarez-Zepeda and D. E. Martire (Washington, DC, USA) . . . . . 285

Stationary phase effects in reversed-phase liquid chromatography under overload conditions
by N. T. Miller (Conshohocken, PA, USA) . . . . . 301

Structure–retention correlation in liquid chromatography (Review)
by T. Hanai (Kyoto, Japan) . . . . . 313

Retention prediction for polymer additives in reversed-phase liquid chromatography
by K. Jinno and Y. Yokoyama (Toyohashi, Japan) . . . . . 325

Retention prediction of analytes in reversed-phase high-performance liquid chromatography based on molecular structure. VI. Disubstituted aromatic compounds
by R. M. Smith and C. M. Burr (Loughborough, UK) . . . . . 335

*Mobile and stationary phases*

Chiral stationary phases derived from tyrosine (Review)
by M. Caude (Paris, France), A. Tambuté (Vert-le-Petit, France) and L. Siret (Paris, France) 357

High-performance liquid chromatography of amino acids, peptides and proteins. CIX. Investigations on the relation between the ligand density of Cibacron Blue immobilized porous and non-porous sorbents and protein-binding capacities and association constants
by H. J. Wirth, K. K. Unger and M. T. W. Hearn (Clayton, Australia) . . . . . 383

Multivariate characterization of solvent strength and solvent selectivity in reversed-phase high-performance liquid chromatography
by P. M. J. Coenegracht, A. K. Smilde, H. Benak, C. H. P. Bruins, H. J. Metting, H. de Vries and D. A. Doornbos (Groningen, Netherlands) . . . . . 397

Dependence of retention on the organic modifier concentration and multicomponent adsorption behavior in reversed-phase chromatography
by F. D. Antia and Cs. Horváth (New Haven, CT, USA) . . . . . 411

Gradient elution methods for predicting isocratic conditions (Review)
by P. J. Schoenmakers (Eindhoven, Netherlands), Å. Bartha (Södertälje, Sweden) and H. A. H. Billiet (Delft, Netherlands) . . . . . 425

*Detection*

Standardization of a multi-wavelength UV detector for liquid chromatography-based toxicological analysis
by S. R. Binder, A. K. Adams, M. Regalia and H. Essien (Hercules, CA, USA) and R. Rosenblum (Rehovot, Israel) . . . . . 449

Gas chromatographic detectors for use in column liquid chromatography (Review)
by Ch. E. Kientz (Rijswijk, Netherlands), G. J. de Jong (Weesp, Netherlands) and U. A. Th. Brinkman (Amsterdam, Netherlands) . . . . . 461

Possibilities of enhancing the sensitivity of the determination of UV-absorbing compounds in high-performance liquid chromatography
by P. Jandera and B. Prokeš (Pardubice, Czechoslovakia) . . . . . 495

*Applications*

Chromatographic characterization of some dideoxyribonucleosides
by N.-I. Jang and P. R. Brown (Kingston, RI, USA) and E. Grushka (Jerusalem, Israel) . 507

Chromatographic separation and examination of carbohydrate and phenolic components of the non-tannin fraction of black wattle (*Acacia mearnsii*) bark extract
by S. C. Churms and A. M. Stephen (Cape Town, South Africa) . . . . . 519

Multi-step procedure for the separation of vinyl chloride oligomers
by J. V. Dawkins and M. J. Forrest (Loughborough, UK) and M. J. Shepherd (Norwich, UK) . . . . . 539

Separation and detection of oxidation products in Neurolite raw material
by P. A. Ryan, B. A. Ewels and J. L. Glajch (N. Billerica, MA, USA) . . . . . 549

Determination of free cyanide in gold cyanidation process liquors by ion-interaction chromatography with post-column derivatization
by P. A. Fagan and P. R. Haddad (Kensington, Australia) . . . . . 559

Chromatographic data for pharmacological classification of imidazol(in)e drugs
by R. Gami-Yilinkou and R. Kaliszan (Gdańsk, Poland) . . . . . 573

Use of Ce(IV) oxidizing agent for the derivatization of polycyclic aromatic hydrocarbons for liquid chromatography–electrochemical detection
by J. R. Mazzeo and I. S. Krull (Boston, MA, USA) and P. T. Kissinger (West Lafayette, IN, USA) . . . . . 585

Confirmation of domoic acid in shellfish using butyl isothiocyanate and reversed-phase liquid chromatography
by J. F. Lawrence and C. Ménard (Ottawa, Canada) . . . . . 595

Determination of hydroxy and peroxy acid derivatives of uroporphyrin in the plasma of patients with congential erythropoietic prophyria by high-performance liquid chromatography
by R. Guo and C. K. Lim (Harrow, UK) . . . . . 603

Chemical class separation of coal liquids by high-performance liquid chromatography on a semi-preparative, phosphoric acid-modified amino-bonded-phase column
by J. W. Haas, III and P. C. Uden (Amherst, MA, USA) . . . . . 609

Chiral separation of ibutilide enantiomers by derivatization with 1-naphthyl isocyanate and high-performance liquid chromatography on a Pirkle column
by C. L. Hsu and R. R. Walters (Kalamazoo, MI, USA) . . . . . 621

Separation of copolymers according to composition with special emphasis on the effect of block structure
by G. Glöckner (Dresden, Germany) and J. H. M. van den Berg (Weesp, Netherlands) . . 629

Analysis of steroids. XLI. Ion-pair high-performance liquid chromatographic separation of quaternary ammonium steroids on silica
by M. Gazdag, M. Babják, P. Kemenes-Bakos and S. Görög (Budapest, Hungary) . . . 639

THIN-LAYER CHROMATOGRAPHY

Mechanism of liquid adsorption chromatography in thin-layer chromatography with ternary mobile phases
by B. Ościk-Mendyk (Lublin, Poland) . . . . . 645

Computer-assisted sample clean-up in liquid chromatography from thin-layer chromatographic data
by A. M. Siouffi and N. Volpe (Marseille, France) . . . . . 657

Comparative investigation of the retention behaviour of nucleoside derivatives on alumina stationary phases in thin-layer chromatography and high-performance liquid chromatography
by K. Valkó, T. Cserháti and E. Forgács (Budapest, Hungary) . . . . . 667

COUNTER-CURRENT CHROMATOGRAPHY

Practical approach to high-speed counter-current chromatography
by A. Berthod (Villeurbanne, France) . . . . . 677

Protein separation with aqueous–aqueous polymer systems by two types of counter-current chromatographs
by Y. Shibusawa and Y. Ito (Bethesda, MD, USA) . . . . . 695

SIZE-EXCLUSION CHROMATOGRAPHY

Effect of sample viscosity in high-performance size-exclusion chromatography and its control
by M. Czok, A. M. Katti and G. Guiochon (Knoxville and Oak Ridge, TN, USA) . . . 705

Hydrodynamic and size-exclusion chromatography of polymers on porous particles
by G. Stegeman, J. C. Kraak and H. Poppe (Amsterdam, Netherlands) . . . . . 721

AFFINITY CHROMATOGRAPHY

Isolation of an interleukin 2-binding receptor from activated lymphocytes by high-performance immunoaffinity chromatography
by T. M. Phillips (Washington, DC, USA) . . . . . 741

Cascade-mode multiaffinity chromatography. Fractionation of human serum proteins
by J. Porath and P. Hansen (Uppsala, Sweden) . . . . . 751

SUPERCRITICAL FLUID CHROMATOGRAPHY

Separation of triacylglycerols by supercritical-fluid argentation chromatography
by M. Demirbüker and L. G. Blomberg (Stockholm, Sweden) . . . . . 765

Sorption isotherms in supercritical fluid chromatography (Review)
by C. R. Yonker and R. D. Smith (Richland, WA, USA) . . . . . 775

FIELD-FLOW FRACTIONATION

Comparison of power and exponential field programming in field-flow fractionation
by P. S. Williams and J. C. Giddings (Salt Lake City, UT, USA) . . . . . 787

Quantitative particle-size distributions by sedimentation field-flow fractionation with densimeter detector
by J. J. Kirkland and W. W. Yau (Wilmington, DE, USA) . . . . . 799

ELECTROPHORESIS

High-performance displacement electrophoresis in 0.025- to 0.050-mm capillaries coated with a polymer to suppress adsorption and electroendoosmosis
by S. Hjertén and M. Kiessling-Johansson (Uppsala, Sweden) . . . . . 811

High-performance capillary electrophoresis of proteins. Sodium dodecyl sulphate–polyacrylamide gel-filled capillary column for the determination of recombinant biotechnology-derived proteins
by K. Tsuji (Kalamazoo, MI, USA) . . . . . 823

Axial-beam absorbance detection for capillary electrophoresis
by J. A. Taylor and E. S. Yeung (Ames, IA, USA) . . . . . 831

*Author Index* . . . . . 839

In articles with more than one author, the name of the author to whom correspondence should be addressed is indicated in the article heading by a 6-pointed asterisk (*)